

# Can plasma exchange therapy induce regulatory T lymphocytes in multiple sclerosis patients?

A. Jamshidian and M. Gharagozloo  
Department of Immunology, School of Medicine,  
Isfahan University of Medical Sciences, Isfahan,  
Iran

Accepted for publication 8 December 2011  
Correspondence: M. Gharagozloo, Department  
of Immunology, School of Medicine, Isfahan  
University of Medical Sciences, Isfahan,  
81744-176, Iran.  
E-mail: gharagozlo@med.mui.ac.ir

## Summary

Plasma exchange is used increasingly as an individual therapeutic decision for treating of severe, steroid-resistant relapses of multiple sclerosis (MS). However, given that its mechanism of action in this CD4<sup>+</sup> T cell-mediated autoimmune disease remains unknown, it is not yet considered as a routine therapy for this prevalent neuroimmune disorder. In this regard, we hypothesized that plasma exchange, by depleting the body of inflammatory mediators that acts as providers of co-stimulatory signals for the adaptive immune system, provides the immune system with an exceptional break for *de-novo* recognition of autoantigens in a tolerogenic manner. This may lead to an increase in the frequency and function of myelin-specific regulatory T cells. For evaluating this we suggest some *in vitro* and *in vivo* studies to analyse the effects of varied dilutions of normal and MS plasmas on the induction of regulatory T cells or on the function of isolated and purified regulatory T cells. Clarifying the effects of therapeutic plasma exchange on regulatory T cells as the major controllers of autoimmune responses may provide us with strong evidence to use this procedure as a disease-modifying treatment in remission phase for reducing the rate and severity of future attacks, in addition to more trustworthy therapy in severe relapses of MS.

**Keywords:** multiple sclerosis, plasma exchange, regulatory T cell

## Introduction

Administration of high doses of intravenous corticosteroids (CS), such as methylprednisolone, is the regular therapy for acute relapses of multiple sclerosis (MS). Although this routine treatment is generally effective for relieving symptoms of an attack in the short term, it does not appear to have a momentous impact on long-term recovery from disease. Additionally, in some patients suffering from severe steroid-resistant relapses, the clinical response to CS therapy may be inadequate. Such patients could obtain clinical benefit from following plasma exchange (PE) therapy. PE is considered increasingly as an individual treatment in patients with severe relapses not responding appropriately to CS. However, because of the lack of proper studies and the unknown mechanism of action, PE is not carried out as an enduring disease-modifying strategy in MS patients [1], while it is an established therapy for the treatment of many other neuroimmune disorders, such as myasthenia gravis, Lambert-Eaton syndrome, Guillain-Barré syndrome and chronic demyelinating polyneuropathy [2,3]. The purpose of the PE procedure is to deplete the blood of various immunological

factors, such as antibodies, complement components, chemokines, cytokines and almost all known and unknown inflammatory factors with which their over-production in the patient's body may be associated with the initiation or progression of the disease process [1]. Clearly, this mechanism cannot directly cause all the improving effects of plasma exchange in a cell-mediated autoimmune disease such as MS, in view of the fact that these soluble factors are made by cells and the original producer cells will still be in the body after plasmapheresis is completed. In this regard, some previous studies have shown that plasmapheresis, in addition to depleting the antibodies and other soluble factors from the blood, may affect the frequency and phenotype of different immune cell populations [4–10], and also the function of these cells and production of some cytokines and other soluble factors [11–16]. Regulatory T cells are known as the most important players in maintaining immune system homeostasis and tolerance to self-antigens [17,18], and are also accepted as the main controllers of MS [19–21]. One study has shown that the plasmapheresis treatment of systemic lupus erythematosus (SLE) patients induced a significant increase in the number of peripheral

CD4<sup>+</sup>CD25<sup>(high)</sup>forkhead box protein 2 (FoxP3<sup>+</sup>) regulatory T cells in parallel to the decrease in disease activity [22]. According to these studies, it seems reasonable to suppose that one mechanism by which plasmapheresis exerts its effect on CD4<sup>+</sup> T cell-mediated autoimmune diseases such as MS may be through exertion of some effects on the frequency, ratio and function of regulatory T cells.

### The hypothesis

By depleting the body of inflammatory factors, plasmapheresis may give the immune system an opportunity to re-adjust itself. This may be possible in part via induction of regulatory T cells through reducing the inflammatory cytokines found in the patients' plasma. According to the established 'two-signal' theory in basic immunology, it has long been accepted that immune activation of T cells is irrefutably dependent upon receiving co-stimulatory signals as the second signal in addition to specific antigen recognition by the T cell receptor complex as the primary stimulatory signal [23]. The B7/CD28 ligand/receptor system is one of the most important co-stimulatory pathways. Any disruption occurring in this signalling pathway results in suppression of the immune responses by induction of functional unresponsiveness or anergy in stimulated T lymphocytes, and in some cases brings about antigen-specific tolerance [24]. Soluble inflammatory factors and cytokines of the innate immune system are the most important regulators of the expression of co-stimulatory molecules, such as B7 family members, on the surface of antigen-presenting cells (APCs) [24–26]. Thus, in the presence of inflammation, antigen recognition by specific T cells is often associated with immune activation, whereas in the absence of inflammatory factors the same antigen recognition can cause the induction of tolerance and unresponsiveness in these antigen-specific T cells and confer upon them the phenotype and function of regulatory T cells. Because plasma exchange washes the body of inflammatory agents, or at least dilutes them in the blood and therefore in the whole body, it may provide naive autoreactive T lymphocytes with the opportunity of recognizing related self-antigens without adequate co-stimulatory signals in a tolerogenic *de-novo* manner. This may cause an increase in the induction of inducible subpopulations of myelin-specific regulatory T cells and also some improvement in the function of naturally occurring regulatory T cells and pre-existing inducible regulatory T cells because of the changes in their cytokine milieu.

### Evaluating the hypothesis

For investigating these probable effects, we suggest some *in vitro* and *in vivo* studies, as follows:

1. *In vitro* studies to examine the effects of varied dilutions of plasma or serum on the phenotype and inhibitory functions of regulatory T cells;

2. To examine the effects of MS patients' plasma or serum on the phenotype and regulatory function of these cells in comparison with plasma or serum of healthy individuals; and
3. *In vivo* studies comparing the relative frequency and inhibitory function of these cells before and after plasma exchange therapy course in animal models or in MS patients.

### Implications of the hypothesis

If sufficient scientific evidence is provided to confirm positive effects of plasmapheresis on the frequency and function of regulatory T cells, this treatment will have the potential for reducing the rate and severity of future relapses in MS patients, in addition to relieving the symptoms of a present attack. Also, if this hypothesis is proven, continuing efforts can be proposed for specified filtering of the blood and depleting the body of defined inflammatory factors that contrast with the induction or function of these cells, instead of replacing the whole plasma, to perform a more effective and safe treatment for MS and similar autoimmune disorders.

### Disclosure

The authors declare that they have no competing interests.

### References

- 1 Tumani H. Corticosteroids and plasma exchange in multiple sclerosis. *J Neurol* 2008; **255**:36–42.
- 2 Kaynar L, Altuntas F, Aydogdu I *et al.* Therapeutic plasma exchange in patients with neurologic diseases: retrospective multicenter study. *Transfus Apher Sci* 2008; **38**:109–15.
- 3 Szczepiorkowski ZM, Bandarenko N, Kim HC *et al.* Guidelines on the use of therapeutic apheresis in clinical practice: evidence-based approach from the Apheresis Applications Committee of the American Society for Apheresis. *J Clin Apheresis* 2007; **22**:106–75.
- 4 Dau PC. Increased proliferation of blood mononuclear cells after plasmapheresis treatment of patients with demyelinating disease. *J Neuroimmunol* 1990; **30**:15–21.
- 5 Dau PC, Callahan JP. Immune modulation during treatment of systemic sclerosis with plasmapheresis and immunosuppressive drugs. *Clin Immunol Immunopathol* 1994; **70**:159–65.
- 6 De Luca G, Lugesesi A, Iarlori C *et al.* Prednisone and plasma exchange improve suppressor cell function in chronic inflammatory demyelinating polyneuropathy. *J Neuroimmunol* 1999; **95**:190–4.
- 7 Goto H, Matsuo H, Nakane S *et al.* Plasmapheresis affects T helper type-1/T helper type-2 balance of circulating peripheral lymphocytes. *Ther Apher* 2001; **5**:494–6.
- 8 Kambara C, Matsuo H, Fukudome T, Goto H, Shibuya N. Miller Fisher syndrome and plasmapheresis. *Ther Apher* 2002; **6**:450–3.
- 9 Yoshii F, Shinohara Y. Lymphocyte subset proportions in Guillain-Barré syndrome patients treated with plasmapheresis. *Eur Neurol* 2000; **44**:162–7.

- 10 Yeh JH, Chien PJ, Hsueh YM, Shih CM, Chiu HC. Changes in the lymphocyte subset after double-filtration plasmapheresis. *Am J Clin Pathol* 2007; **128**:940–4.
- 11 Steven MM, Tanner AR, Holdstock GE *et al*. The effect of plasma exchange on the *in-vitro* monocyte function of patients with immune complex diseases. *Clin Exp Immunol* 1981; **45**:240–5.
- 12 Shariatmadar S, Nassiri M, Vincek V. Effect of plasma exchange on cytokines measured by multianalyte bead array in thrombotic thrombocytopenic purpura. *Am J Hematol* 2005; **79**:83–8.
- 13 Dau PC. Increased antibody production in peripheral blood mononuclear cells after plasma exchange therapy in multiple sclerosis. *J Neuroimmunol* 1995; **62**:197–200.
- 14 Yeh JH, Wang SH, Chien PJ, Shih CM, Chiu HC. Changes in serum cytokine levels during plasmapheresis in patients with myasthenia gravis. *Eur J Neurol* 2009; **16**:1318–22.
- 15 Tesar V, Jelinkova E, Jirsa M Jr, Bakosova M, Pitha P, Chabova V. Soluble adhesion molecules and cytokines in patients with myasthenia gravis treated by plasma exchange. *Blood Purif* 2000; **18**:115–20.
- 16 Tesar V, Jelinkova E, Masek Z *et al*. Influence of plasma exchange on serum levels of cytokines and adhesion molecules in ANCA-positive renal vasculitis. *Blood Purif* 1998; **16**:72–80.
- 17 Sakaguchi S. Regulatory T cells: key controllers of immunologic self-tolerance. *Cell* 2000; **101**:455–8.
- 18 Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains [CD25]. Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* 1995; **155**:1151–64.
- 19 Viglietta V, Baecher-Allan C, Weiner HL, Hafler DA. Loss of functional suppression by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. *J Exp Med* 2004; **199**:971–9.
- 20 Venken K, Hellings N, Thewissen M *et al*. Compromised CD4+CD25[high] regulatory T-cell function in patients with relapsing–remitting multiple sclerosis is correlated with a reduced frequency of FOXP3-positive cells and reduced FOXP3 expression at the single-cell level. *Immunology* 2008; **123**:79–89.
- 21 Haas J, Hug A, Viehaver A *et al*. Reduced suppressive effect of CD4+CD25high regulatory T cells on the T cell immune response against myelin oligodendrocyte glycoprotein in patients with multiple sclerosis. *Eur J Immunol* 2005; **35**:3343–52.
- 22 Barath S, Soltesz P, Kiss E *et al*. The severity of systemic lupus erythematosus negatively correlates with the increasing number of CD4+CD25[high]FoxP3+ regulatory T cells during repeated plasmapheresis treatments of patients. *Autoimmunity* 2007; **40**:521–8.
- 23 Baxter A, Hodgkin P. Activation rules: the two-signal theories of immune activation. *Nat Rev Immunol* 2002; **2**:439–46.
- 24 Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Annu Rev Immunol* 1996; **14**:233–58.
- 25 Chang CH, Furue M, Tamaki K. B7-1 expression of Langerhans cells is up-regulated by proinflammatory cytokines, and is down-regulated by interferon-gamma or by interleukin-10. *Eur J Immunol* 1995; **25**:394–8.
- 26 Freedman AS, Freeman GJ, Rhynhart K, Nadler LM. Selective induction of B7/BB-1 on interferon-gamma stimulated monocytes: a potential mechanism for amplification of T cell activation through the CD28 pathway. *Cell Immunol* 1991; **137**:429–37.